

NHS Kernow CCG Joint Dressings Formulary - To be used in conjunction with the order form

Link to Cornwall Joint Formulary ; - <https://www.eclipsesolutions.org/cornwall/>

Cornwall Community Tissue Viability team:- tissue.viabilitycornwall@nhs.net Tel 01208 834371

Heather Newton (heathernewton@nhs.net) 01872 252673

Podiatry: 01209318093 or liza.downing@nhs.net

Cornwall Community (Primary Care and CFT)

NECROTIC		
Hydrocolloid	DuoDERM Extra Thin	Dry to lightly exuding wounds. Wear time 3-7days. Warm before application.
	Comfeel Plus Transparent Comfeel Plus	Comfeel is for low to medium exudate wounds. Can be left in place for a week. Warm before application. Do not use on infected wounds
	ActivHeal Hydrocolloid	Low to moderate level of exudate. Wear time 3-7 days.
Hydrogel	ActivHeal Hydrogel	Use to hydrate dry wounds
	Intrasite Conformable	FOR WOUND DEBRIDEMENT ONLY. Single Use Application Daily. 10 x 10cm (equiv to 8g) 10 x 20cm (equiv to 15g)
	ActiFormCool Sheet	To remain in place for 2-3 days. Use for rehydration and debridement. Cut to size, can be used layered. NOT for cavity wounds
INFECTED/COLONISED		
Antimicrobial Primary Silver Dressing	Aquacel Ag+ Extra	Broad-spectrum antimicrobial and anti-biofilm. For use on dry-high exuding wounds. Can be left in place up to 7 days. Use 2-3 weeks then wound reviewed. Please ensure "+" is included on dressing request.
Alginate/Manuka Honey	Algivon Plus	Alginate dressing impregnated with Manuka Honey
	Activon Tube Activon Tulle	100% Manuka Honey. Gauze dressings impregnated with manuka honey.
	Actilite	Viscose net dressing coated with 99% Manuka Honey and 1% Manuka oil
Antibacterial	Flamazine(Silver Sulfadiazine 1% cream) 50g	Not to be used 1st line for burns. Obtain on FP10. Only use on dry wounds as can increase level of exudate. Apply daily. Max 7 day open storage time.
	Flaminal Hydro (for drier wounds) Flaminal Forte (for wet wounds)	Used only on advice of TV team
Cadaxomer iodine	Iodoflex	
Iodine Tulle	Povitulle Inadine	Arterial, diabetic and trauma wounds. Change when white or wet. Short term single use only. Only via NHS Supply chain
SLOUGHY		
Alginate	Sorbsan Flat Sheet Sorbsan Plus Pad	
	Aquacel Extra Urgoclean Pad	Wear time- 3-7 days depending on exudate (RCHT First line) - For low to moderate wounds.
Rapid Capillary dressing	Vacutex	
Cavity	Sorbsan Ribbon Urgoclean Rope	Use if daily dressings required First line for cavity wounds
Biotherapy	Larvae - Maggots	RCHT order from pharmacy. Cornwall Community - Wound debridement available on FP10. Consult Tissue Viability prior to ordering. Order direct from BioMonde (Tel 0845 2301810)
EPITHELIALISING		
Semi Permeable Adhesive	Hydrofilm	Use as secondary dressing to aid debridement or to protect newly epithelializing wounds. Used to reduce friction.
Adhesive Island	Hydrofilm Plus Cosmopore	Waterproof / bacteria proof surgical wound dressing Not waterproof NB RCHT only for awkward areas
ODOUR		
Charcoal Dressing (Non Absorbent)	CliniSorb	Carbon dressing for malodorous wound.
	Anabact	Obtain on FP10. Metronidazole gel (POM) for malodorous wounds.
GRANULATING		
Foam Adhesive	UrgoTul Absorb Border Aquacel Foam Adhesive	Indicated for the treatment of low to moderately exuding wounds Multi-layered absorbent foam dressing for mid-high exuding wounds with a silicone adhesive border. Wound contact layer containing Aquacel
	Allevyn life	Indicated for the treatment of high exuding wounds. Used only on advice from TV team
Foam Non-adhesive	UrgoTul Absorb	Indicated for the treatment of low to moderate exuding wounds
	Aquacel Foam Non-Adhesive	Multilayered absorbent foam dressing for mid-high exuding wounds. Wound contact layer containing Aquacel.

NHS Kernow CCG Joint Dressings Formulary - To be used in conjunction with the order form

Link to Cornwall Joint Formulary ; - <https://www.eclipsesolutions.org/cornwall/>

Cornwall Community Tissue Viability team:- tissue.viabilitycornwall@nhs.net Tel 01208 834371

Heather Newton (heathernewton@nhs.net) 01872 252673

Podiatry: 01209318093 or liza.downing@nhs.net

Non-adherent	Atrauman	Use where contact layer requires changing 3-4 times weekly
Silicone	Adaptic Touch	Use where dressing needs to stay in place for 7 – 14 days. Renew outer bandages / dressings as required
Interactive	Urgostart Range <ul style="list-style-type: none"> • Urgostart Plus Pad • Urgostart Plus Border • Urgostart Contact 	Protease inhibitor action for Diabetic Foot Ulcer (DFU) and Venous Leg Ulcer (VLU) on Tissue Viability advice only
MISCELLANEOUS		
Barrier	Medi Derma-S Barrier Film 1ml and 3ml	Prevents maceration and for general skin protection. Ensure correct application. One application stick lasts 72 hours. DO NOT OVER APPLY.
	Medi Derma-S Barrier cream 28g	28g tube should be sufficient for 1 months' supply. Pea-sized amount applied daily or every 3rd to 4th episode of incontinence.
Absorbent Cellulose Dressing	Zetuvit Sterile Zetuvit Plus	Absorbent cellulose dressing with fluid repellent backing, use for low level exudate Absorbent cellulose dressing with fluid repellent backing, use for heavy exudate
Topical Negative Pressure Therapy	VAC/ActiVAC	
	PICO 7	Dressings available on FP10. Consists of 1 pump, 2 sterile dressings, sterile retention strips and 2 AA batteries. Use in the community is following TV recommendation only. For use on chronic wounds (i.e. > 6 weeks) only. Wound reviewed by TV to assess response to treatment. Discontinue if wound becomes static.
Wound Irrigation	Clinipod Irripod	Irrigate only if loose debris present.
Surgical Tape	Scanpore	Dressing and bandage retention
Bleeding wounds	Kaltostat	To aid the cessation of bleeding in wounds
Keloid Scarring	Cica-Care	Tissue Viability Team advice
Dressing Pack	Nurse-It	Order M/L glove size
	Richardson Wound care pack (option 11)	Direct purchase only
Gauze Swabs	Sterile (7.5cm)	Pack of 5 gauze swabs
	Non-sterile	Pack of 100 gauze swabs
Debridement	UCS Cloth - for skin care of lower limbs	
Tracheostomy	Trachi-Dress F	For use as directed by interventional radiology team for the management of weekly nephrostomy dressing changes (used in conjunction with waterproof Hydrofilm cover)
BANDAGES		
Type 1 retention bandage	K-band	
Crepe Bandage	Hospicrepe	
Multilayer compression	UrgoKTwo	
Cohesive Short stretch	Actico	
Stockinette	Acti-Fast blue-7.5cm Acti-Fast yellow-10.75cm	
Elasticated tubular bandage	easiGRIP	
Paste bandage	Viscopaste Zipzoc	
Padding	K-Soft	
	Profore #1	For use only when allergic to K-Soft
Compression Hosiery	Medi Range	
Lymphoedema Garments	Juxta Wrap	Specialist only - Not direct purchase