

NHS Kernow CCG Joint Dressings Formulary - To be used in conjunction with the order form

Link to Cornwall Joint Formulary ; - <https://www.eclipsesolutions.org/cornwall/>

Cornwall Community Tissue Viability team:- tissue.viabilitycornwall@nhs.net Tel 01208 834371

Heather Newton (heathernewton@nhs.net) 01872 252673

Podiatry: 01209318093 or liza.downing@nhs.net

Wound MANAGEMENT- RCHT

NECROTIC		
Hydrocolloid	Comfeel Plus Transparent Comfeel Plus	Comfeel Plus Transparent is for low exudate wounds. Can be left in place for a week. Warm before application. Comfeel Plus can be used on wounds with moderate exudate Do not use on infected wounds
Hydrogel	ActivHeal Hydrogel	Use to hydrate dry wounds
	ActiFormCool Sheet	To remain in place for 2-3 days. Use for rehydration and debridement. Cut to size. NOT for cavity wounds
INFECTED/COLONISED		
Antimicrobial Primary Silver Dressing	Aquacel Ag+ Extra (Equipment Library Only)	Broad-spectrum antimicrobial and anti-biofilm. Can be used on wet wounds. Can be left in place up to 7 days. Use 2-3 weeks then wound reviewed.
	Urgotul Silver	To be used under NPWT and on low exudating wounds
Alginate/Manuka Honey	Algivon Plus	Alginate dressing impregnated with Manuka Honey (Obtain from RCHT Equipment Library)
	Activon Tube Activon Tulle	Ointment and dressings impregnated with Manuka honey. (Obtain from RCHT Equipment Library)
Antibacterial	Flamazine (Silver Sulfadiazine 1% cream) 50g	Not to be used 1st line for burns. Can increase level of exudate. Apply daily. Max 7 day open storage time.
	Flaminal Hydro (blue)POM Flaminal Forte (yellow)POM	For drier wounds - Used only on advice of TV team For wet wounds - Used only on advice of TV team
Cadaxomer iodine	Iodoflex	For use on wet wounds (Obtain from pharmacy)
Iodine Tulle	Inadine	Arterial, diabetic and trauma wounds. Not for use on wet wounds Change when white or wet. Short term single use only.
SLOUGHY		
Protease Modulating Matrix	Urgoclean Pad	For low to moderate wounds.
Cavity	Urgoclean Rope	First line for cavity wounds
Biotherapy	Larvae - Maggots	RCHT order from pharmacy.
EPITHELIALISING		
Semi Permeable Adhesive	Opsite Flexigrid	Use as secondary dressing to aid debridement or to protect newly epithelializing wounds. Used to reduce friction.
Adhesive Island	Opsite Post Op Cosmopore	Waterproof / bacteria proof surgical wound dressing Not waterproof Not first line for surgical wounds due to lack of bacteria and waterproofing capacity
ODOUR		
Charcoal Dressing (Non Absorbent)	CliniSorb	Carbon dressing for malodorous wound.
GRANULATING		
Silicone Foam Adhesive	UrgoTul Absorb Border	Low to moderately exuding wounds. Primary dressig First line skin tear dressing- not to be applied daily maximum 3-4 dressings per week
Silicone foam	UrgoTul Absorb	
Non-adherent	Atrauman	Use where contact layer requires changing 3-4 times weekly
Silicone	Adaptic Touch	Use where dressing needs to stay in place for 7 – 14 days. Renew outer bandages / dressings as required
MISCELLANEOUS		
Barrier	Medi Derma-S Barrier Film 1ml and 3ml	Prevents maceration and for general skin protection. Ensure correct application. One application stick lasts 72 hours. DO NOT OVER APPLY.
	Medi Derma-S Barrier cream 28g	28g tube should be sufficient for 1 months' supply. Pea-sized amount applied daily or every 3rd to 4th episode of incontinence.
Absorbent Cellulose Dressing	Zetuvit Plus Eclipse Range	First line – superabsorbent secondary dressing for moderate to heavily exuding wounds Superabsorbent secondary dressing for heavily exuding wounds
Topical Negative Pressure Therapy	PICO 7	Available via RCHT Equipment library
Bleeding wounds	Kaltostat	To aid the cessation of bleeding in wounds
Type 1 retention bandage	K-band	
Crepe Bandage	Hospicrepe	
Multilayer compression	UrgoKTwo	
Cohesive Short stretch	Actico	
Stockinette	Clini-Fast blue-7.5cm Clini-Fast yellow-10.75cm	
Paste bandage	Zipzoc	
Padding	K-Soft	
Compression Hosiery	Altiform	