

## NHS Kernow CCG Joint Dressings Formulary - To be used in conjunction with the order form

Link to Cornwall Joint Formulary ; - <https://www.eclipsesolutions.org/cornwall/>

Cornwall Community Tissue Viability team:- [tissue.viabilitycornwall@nhs.net](mailto:tissue.viabilitycornwall@nhs.net) Tel 01208 834371

Heather Newton ([heathernewton@nhs.net](mailto:heathernewton@nhs.net)) 01872 252673

Podiatry: 01872 221435/221436 or [liza.downing@nhs.net](mailto:liza.downing@nhs.net)/[neil.smith29@nhs.net](mailto:neil.smith29@nhs.net)

## Wound MANAGEMENT- PODIATRY

<b>NECROTIC</b>		
Hydrogel	ActivHeal Hydrogel	Use to hydrate dry wounds
	ActiFormCool Sheet	To remain in place for 2-3 days. Use for rehydration and debridement. Cut to size, can be used layered. NOT for cavity wounds
<b>INFECTED/COLONISED</b>		
Antimicrobial Primary Silver Dressing	Urgotul Silver	Effective against MRSA and Pseudomonas spp. Urgotul Silver can be left in place for up to a week. Max use 2-3 weeks then review use
Antibacterial	Flaminal Hydro (blue) Flaminal Forte (yellow)	For dry wounds For wet wounds
Cadaxomer iodine	Iodoflex	
Iodine Tulle	Inadine	Arterial, diabetic and trauma wounds. Change when white or wet. Short term single use only. <b>Only via NHS Supply chain</b>
<b>SLOUGHY</b>		
Protease Modulating Matrix	Aquacel Extra	Wear time- 3-7 days depending on exudate
Protease Inhibitor	Urgostart Plus	Poly absorbent fibre pad containing TLC-NOSF. For all wound healing stages. Do not use on bleeding wounds
Alginate Flat Sheet	Sorbsan Flat	Use where daily applications are required due to wound site requiring daily bathing
Alginate Pad	Sorbsan Plus	As above. Alginate with absorbent pad
Biotherapy	Larvae - Maggots	<b>RCHT order from pharmacy.</b>
<b>GRANULATING</b>		
Non-adherent	Atrauman	Use where contact layer requires changing 3-4 times weekly
	Telfa	Absorbent, perforated plastic film faced dressing
Protease Inhibitor	Urgostart	Soft absorbent foam containing protease inhibitor (TLC-NOSF). NICE evidence supports use to treat DFUs. Do not use infected / critically colonised wounds
Non Adhesive Foam	Advasorb	Hydrophilic foam. Low to highly exuding wounds. Do not use on bleeding or arterial wounds
	Urgotul Absorb Non-Adhesive	First line skin tear dressing. Not to be applied daily, maximum 3-4 dressings per week.
Soft Adhesive Foam	Biatain Soft Hold	Light to moderate exudates
<b>EPITHELIALISING</b>		
Adhesive Island	Cosmopore	Not waterproof
<b>MISCELLANEOUS</b>		
Barrier	Medi Derma-S Barrier Film 1ml and 3ml	Prevents maceration and for general skin protection. Ensure correct application. One application stick lasts 72 hours. DO NOT OVER APPLY.
Topical Negative Pressure Therapy	VAC	Diabetic Specialist Podiatrist use only
Wound Irrigation	Clinipod Irripod	
Surgical Tape	Mefix	Dressing and bandage retention
Bleeding wounds	Kaltostat	To aid the cessation of bleeding in wounds – nail surgery use only
Keloid Scarring	Cica-Care	Ordered via Supply Chain
Contact Layer	Jelonet Paraffin Dressing	For use in nail surgery
<b>BANDAGES</b>		
Crepe Bandage	Hospicrepe	
Stockinette	Acti-Fast blue-7.5cm	
	Acti-Fast yellow-10.75cm	
Padding	K-Soft	
	Profore #1	For use only when allergic to K-Soft