

## Cornwall Community (Primary Care and CFT)

Link to Cornwall Formulary: - <https://www.eclipsesolutions.org/cornwall/>

Cornwall Community Tissue Viability team: - [tissue.viabilitycornwall@nhs.net](mailto:tissue.viabilitycornwall@nhs.net) Tel 01208 834371

NHS Cornwall and Isles of Scilly Intergrated Care Board Prescribing: - [ciosicb.prescribing@nhs.net](mailto:ciosicb.prescribing@nhs.net)

Podiatry: 01209 318093

NECROTIC		
Hydrocolloid	Duoderm Extra Thin	Duoderm Extra Thin is for low exudate wounds. It can be left in place for a week if required. Warm before application.
	Duoderm Signal	Duoderm Signal can be used on wounds with moderate exudate. Do not use on infected wound.
Hydrogel	ActivHeal Hydrogel	Use to hydrate dry wounds.

INFECTED/COLONISED		
Antimicrobial Primary Silver Dressing	Aquacel Ag+ Extra	Broad-spectrum antimicrobial and anti-biofilm. Can be used on wet wounds. Can be left in place up to 7 days. Use 2-3 weeks then wound reviewed.
Alginate/Manuka Honey	Algivon Plus	Alginate dressing impregnated with Manuka Honey. Useful for wetter wounds.
	Activon Tube Activon Tulle	Ointment impregnated with Manuka honey. (Obtain from RCHT Equipment Library) Tulle dressing impregnated with Manuka honey. Suitable for shallow wounds and those with lower exudate. Discuss with TV for further advice and most appropriate product.
	Actilite	Viscose net dressing coated with 99% Manuka Honey and 1% Manuka oil.
Antibacterial	Flamazine (Silver Sulfadiazine 1% cream) 50g	Not to be used 1st line for burns. Topical antimicrobial cream. Can increase level of exudate therefore daily application recommended. Max 7-day open storage time. Available on FP10 only.
	Flaminal Hydro (blue) Flaminal Forte (yellow)	Contains alginate and antibacterial enzyme. Used for dry wounds. Contains alginate and antibacterial enzyme. Used for wet wounds.
	Prontosan gel X 30ml	Highly viscous gel cleansing, moisturising, and decontaminating wounds and burns. Apply up to a 3mm layer of Prontosan Gel X to the wound bed.
Cadaxomer iodine	Iodoflex	For use on wet wounds. Do not use on patients with Thyroid disorders or sensitivity to Iodine. (Obtain from pharmacy)

INFECTED/COLONISED Cont.		
Povidone-Iodine	Inadine	Not for use on wet wounds or on patients with sensitivity to Iodine. Colour change indicates when to change dressing. Short term single use only.
Topical antimicrobial irrigation	Prontosan solution 350ml	Apply soaked gauze to the wound and leave in situ for 10 minutes.

SLOUGHY		
Protease Modulating Matrix	Aquacel Extra	Wear time- 3-7 days depending on exudate.
Non-adherent poly absorbent fibre	Urgoclean Pad	For low to moderate wounds.
	Urgoclean Rope	Record the number of ropes inserted into the wound cavity. Leave end of rope visible.
	Sorbsan Ribbon	Use if daily dressings required.
Interactive	Urgostart Plus Pad	For the treatment of lower limb wounds Protease inhibitor action.
	Urgostart Plus Border	Follow Urgostart pathway.
	Urgostart Contact	
Hydro-responsive Wound Dressing	Hydroclean Advance	Suitable to treat dry necrotic to wet sloughy wounds, removes devitalised tissue and contributes significantly to wound bed preparation. Available in cavity variant as well.
Biotherapy	Larvae - Maggots	Wound debridement available on FP10. Consult TV prior to ordering. Order direct from BioMonde. (Tel 0845 2301810)

EPITHELIALISING		
Semi Permeable Adhesive	Hydrofilm	Use as secondary dressing to aid debridement or to protect newly epithelializing wounds. Used to reduce friction.
	Cosmopor Transparent	Waterproof / bacteria proof surgical wound dressing.
	Cosmopor E	Fluid repellent island dressing for minor injuries. Not first line for surgical wounds due to lack of bacteria and waterproofing capacity.

ODOUR		
Charcoal Dressing (Non Absorbent)	CliniSorb	Carbon dressing for malodorous wound.
	Anabact gel	Obtain on FP10. Metronidazole gel (POM) for malodorous wounds.

GRANULATING		
Foam Adhesive	UrgoTul Absorb Border	Low to moderately exuding wounds. First line skin tear dressing - not to be applied daily.
	RespoSorb Silicone Border  (previously known as Zetuvit Plus Silicone Border)	Super Absorbent Polymer dressing with silicone wound contact layer and adhesive border which absorbs exudate and binds and retains bacteria and MMPs. Suitable for all exuding wounds and under compression. Showerproof and bacteria proof with transparent border that can be cut. Can be used as primary or secondary dressing and can stay in place for up to 7 days.
	Aquacel Foam Adhesive	Multi-layered absorbent foam dressing for mid-high exuding wounds with a silicone adhesive border. Wound contact layer containing Aquacel.
Foam Non-adhesive	UrgoTul Absorb	Indicated for the treatment of low to moderate exuding wounds.
	RespoSorb Silicone  (previously known as Zetuvit Plus Silicone)	Super Absorbent Polymer dressing with silicone wound contact layer and water repellent outer layer, suitable for moderate to highly exuding wounds and under compression.
	Aquacel Foam Non-Adhesive	Multilayered absorbent foam dressing for mid-high exuding wounds. Wound contact layer containing Aquacel.
Non-adherent	Atrauman	Use where contact layer requires frequent changing. Petroleum free.
Silicone	Adaptic Touch	Use where dressing needs to stay in place for 7 – 14 days. Renew outer bandages / dressings as required

MISCELLANEOUS		
Barrier	Medi Derma-S Barrier Film 1ml and 3ml	Prevents maceration and for general skin protection. Ensure correct application. One application stick lasts 72 hours. DO NOT OVER APPLY.
	Medi Derma-S Barrier cream 28g	28g tube should be sufficient for 1 months' supply. Pea-sized amount applied daily or every 3rd to 4th episode of incontinence.
	Medi-Derma-S pump spray 30ml	Useful for larger areas of moisture associated skin damage.

MISCELLANEOUS Cont.		
Absorbent Cellulose Dressing	Zetuvit	Absorbent cellulose dressing with fluid repellent backing, use for low level exudate.
	Zetuvit Plus	Superabsorbent dressing that contains a blend of cellulose fluff and fluid-retaining superabsorbent particles. Water-repellent, air permeable, non-woven layer protects against contamination. Soft consistency for a cushioning effect. Suitable for use under compression.
Topical Negative Pressure Therapy	VAC/ActiVAC	Available on tissue viability recommendation. TV will order VAC unit only. Dressings and accessories / consumables are available to order via Amecare by teams providing treatment. Portable negative pressure therapy device.
	PICO 7	Single use portable negative pressure therapy device. Manages low volumes of exudate. Use in the community following TV recommendation only. Wound to be reviewed by TV to assess response to treatment. Discontinue if wound becomes static. Please order via CCOMS.
	SNAP	Single use portable negative pressure therapy device. Manages low volumes of exudate. Use in the community following TV recommendation only. Wound to be reviewed by TV to assess response to treatment. Discontinue if wound becomes static. Please order via CCOMS.
Wound Irrigation	Clinipod	Irrigate only.
Surgical Tape	Scanpore	Dressing and bandage retention.
Bleeding wounds	Kaltostat	To aid the cessation of bleeding in wounds.
Dressing Pack	Dressing Pack	S,M,L sizes available
Gauze Swabs	Sterile (7.5cm) Non-sterile	Pack of 5 gauze swabs. Pack of 100 gauze swabs.
Debridement	UCS Cloth - for skin care of lower limbs	Debridement cloth.

BANDAGES		
Type 1 retention bandage	K-band	Dressing retention bandage and component of three-layer compression system.
Crepe Bandage	Hospiform	Dressing retention bandage and component of three-layer compression system.
Multilayer compression	UrgoKTwo	Full compression system measure ankle to ensure correct size is ordered.
	UrgoKTwo reduced	For patients unable to tolerate full compression measure ankle to ensure correct size is ordered.
Cohesive Short stretch	Actico	Inelastic compression bandage to be use in short stretch compression system.
Stockinette	Acti-Fast blue-7.5cm	Retention and limb protection.
	Acti-Fast yellow-10.75cm	
Elasticated tubular bandage	easiGRIP	Retention and limb protection.
Paste bandage	Steri paste	Medicated bandage for dermatology conditions patch test required. Must be applied in correct fashion, competent staff only to apply.
	Zipzoc	On dermatology recommendation or where steri paste application is not possible.
Padding	K-Soft	Limb protection and component of compression systems.
	Profore #1	For use only when allergic to K-Soft.
Compression Hosiery	Medi Range	Available via direct purchase.
Lymphoedema Garments	Juxta Wrap	Compression wraps available on FP10 following tissue viability advice.